

## Contents

### Special Issue: Ultra-High Pressure Metamorphic Rocks International Lithosphere Programme Contribution 345

Preface . . . . .	ix
High-pressure blueschists and eclogites in Hong'an: a framework for addressing the evolution of high- and ultrahigh-pressure rocks in central China E.A. Eide and J.G. Liou . . . . .	1
Sm–Nd geochronology and petrography of garnet pyroxene granulites in the northern Sulu region of China and their geotectonic implication M. Zhai, B. Cong, J. Guo, W. Liu, Y. Li and Q. Wang . . . . .	23
Petrology of ultrahigh-pressure eclogites from the ZK703 drillhole in the Donghai, eastern China Z. Zhang, Z. Xu and H. Xu . . . . .	35
Experimental synthesis and phase relations of phengitic muscovite from 6.5 to 11 GPa in a calcareous metapelite from the Dabie Mountains, China K.J. Domanik and J.R. Holloway . . . . .	51
Stable isotope geochemistry of marbles from the coesite UHP terrains of Dabieshan and Sulu, China D. Rumble III, Q. Wang and R. Zhang . . . . .	79
Decompression $P$ – $T$ path of coesite eclogite to granulite from Weihai, eastern China S. Banno, M. Enami, T. Hirajima, A. Ishiwatari and Q.C. Wang . . . . .	97
A transitional eclogite- to high pressure granulite-facies overprint on coesite–eclogite at Taohang in the Sulu ultrahigh-pressure terrane, Eastern China Y. Yao, K. Ye, J. Liu, B. Cong and Q. Wang . . . . .	109
Metamorphic evolution, mineral chemistry and thermobarometry of schists and orthogneisses hosting ultra-high pressure eclogites in the Dabieshan of central China D.A. Carswell, R.N. Wilson and M. Zhai . . . . .	121
Large areal extent of ultrahigh-pressure metamorphism in the Sulu ultrahigh-pressure terrane of East China: new implications from coesite and omphacite inclusions in zircon of granitic gneiss K. Ye, Y. Yao, I. Katayama, B. Cong, Q. Wang and S. Maruyama . . . . .	157
Eclogites and eclogites in the Western Gneiss Region, Norwegian Caledonides S.J. Cuthbert, D.A. Carswell, E.J. Krogh-Ravna and A. Wain . . . . .	165
Garnet–omphacite–phengite thermobarometry of eclogites from the coesite-bearing unit of the southern Dora-Maira Massif, Western Alps E.U. Nowlan, H.-P. Schertl and W. Schreyer . . . . .	197
Coesite inclusions and prograde compositional zonation of garnet in whiteschist of the HP-UHPM Kokchetav massif, Kazakhstan: a record of progressive UHP metamorphism C.D. Parkinson . . . . .	215
Petrotectonic evolution of the high- to ultrahigh-pressure Maksyutov Complex, Karayanova area, south Ural Mountains: structural and oxygen isotope constraints M.L. Leech and W.G. Ernst . . . . .	235
A possible UHP-eclogite in the Leaota Mts. (South Carpathians) and its history from high-pressure melting to retrograde inclusion in a subduction melange G. Săbău . . . . .	253
Author index to volume 52 . . . . .	277

# Ultrahigh Pressure

The ultrahigh pressure (UHP) technique is a powerful tool for studying the behavior of materials under conditions of extreme pressure. It is typically achieved by the use of diamond anvils, which are pressed together to create a small, high-pressure region. This region is then used to compress a sample material, allowing researchers to study its properties under conditions that are otherwise inaccessible. The UHP technique has been used to study a wide range of materials, including metals, minerals, and polymers, and has provided valuable insights into their behavior under extreme conditions.

One of the key advantages of the UHP technique is its ability to create pressures that are much higher than those achievable by conventional methods. This allows researchers to study materials under conditions that are closer to those found in the Earth's interior, providing a more realistic picture of their behavior. Additionally, the UHP technique is highly versatile, allowing for the study of a wide range of materials and properties. This makes it a valuable tool for researchers in a variety of fields, including geology, materials science, and physics.

Despite its many advantages, the UHP technique does have some limitations. One of the main challenges is the difficulty of creating and maintaining the high-pressure region. This requires the use of specialized equipment and techniques, which can be expensive and time-consuming. Additionally, the small size of the high-pressure region can limit the amount of material that can be studied, making it difficult to obtain large samples for analysis.

